




# THE RADIATION POWER SPECTRUM.



# TT-TE Power Spectrum.




The radiation power spectrum has very distinctive features:

1. Plateau at low multipoles
2. Peaks of different height
3. Trough of similar height
4. Damping at high multipoles



## Multipoles.



Each Multipole is associated with a distinctive angular scales.



## Angles & Multipoles.

♣ Any spherical harmonic  $Y_{lm}$  has  $2l + 1$  zeros in the azimuthal direction and  $l$  zeros in the polar direction. However, the angular scales in the azimuthal direction vary with latitude. An approximate angular scale can be obtained as:

$$\Omega = \frac{4\pi}{l(2l + 1)} \approx \frac{2\pi}{l^2} \Rightarrow \alpha_l \sim \sqrt{\Omega} = \frac{\sqrt{2\pi}}{l} = \frac{144^\circ}{l}$$



## Physics of the Baryon-Photon Plasma.

- Ma & Bertschinger (1996) ApJ; Hu & Dodelson (2002) ARAA.

- ♣ Prior to recombination baryons and photons are strongly coupled. The effect of baryons is to exchange energy-momentum between photons at different frequencies. We can neglect the effect of gravity.

- ♣ Boltzmann equation shows that the monopole and dipole moments of the temperature field in the previous approximation is:

$$\dot{\Theta}_0 = -\frac{4}{3}\Theta_1 \quad \dot{\Theta}_1 = -\frac{k^2}{3}\Theta_0$$

where derivatives are with respect to conformal time.



## Acoustic Oscillations

♣ Since the sound speed is  $c_s = \sqrt{\dot{p}/\dot{\rho}} = 1/\sqrt{3}$ , both equations can be coupled to give:

$$\ddot{\Theta}_0 + c_s^2 k^2 \Theta_0 = 0$$

Physically, these temperature oscillations represent the heating and cooling of a fluid that is compressed and rarefied by an standing acoustic wave. This behaviour continues until recombination.

♣ The temperature distribution at recombination will be:

$$\Theta(k, \tau_{rec}) = A \cos(k c_s \tau_{rec})$$

In here:  $c_s \tau_{rec}$  is the distance sound can travel by the time  $\tau_{rec}$ , called **sound horizon**.



## Peak Structure.

♣ The radiation power spectrum is the variance of  $\Theta(k, \tau_{rec})$ ; therefore, modes that are caught at maxima or minima of their oscillation at recombination correspond to peaks in the power spectrum. This is the origin of the 'peaks' detected in the observations.

♣ How does this spectrum of inhomogeneities at recombination appear to us today?. A wavelength  $\lambda$  appears as an angular anisotropy of scale  $\theta \approx \lambda/d_A$ , with  $d_A$  the angular diameter distance. Therefore, peaks should appear at

$$l_n \sim n \frac{\pi d_{A,rec}}{c_s \tau_{rec}}$$

Therefore, the position of the acoustic peaks can be used to determine the geometry of the Universe.

♣ At small scales, the coupling of baryons and photons is not so perfect and photons leak out and erase density perturbations (SILK DAMPING).



## Parameter Sensitivity.

♡ If we denote by  $l_a = \pi d_{A,rec}/c_s \tau_{rec}$  the multipole corresponding to the angular extend of the sound horizon, by  $l_{eq} = k_{eq} d_{A,rec}$  the particle horizon at matter-radiation equality and  $l_d$  the Silk damping scale, then for a large class of cosmological models, cosmological parameters are approximately given by:

$$\begin{aligned}\frac{\Delta l_a}{l_a} &\approx -0.24 \frac{\Delta \Omega_m h^2}{\Omega_m h^2} + 0.07 \frac{\Delta \Omega_b h^2}{\Omega_b h^2} - 0.17 \frac{\Delta \Omega_\Lambda}{\Omega_\Lambda} - 1.1 \frac{\Delta \Omega_{tot}}{\Omega_{tot}} \\ \frac{\Delta l_{eq}}{l_{eq}} &\approx 0.5 \frac{\Delta \Omega_m h^2}{\Omega_m h^2} + -0.17 \frac{\Delta \Omega_\Lambda}{\Omega_\Lambda} - 1.1 \frac{\Delta \Omega_{tot}}{\Omega_{tot}} \\ \frac{\Delta l_d}{l_d} &\approx -0.21 \frac{\Delta \Omega_m h^2}{\Omega_m h^2} + 0.20 \frac{\Delta \Omega_b h^2}{\Omega_b h^2} - 0.17 \frac{\Delta \Omega_\Lambda}{\Omega_\Lambda} - 1.1 \frac{\Delta \Omega_{tot}}{\Omega_{tot}}\end{aligned}$$





# CMB TEMPERATURE ANISOTROPIES.



## Sources of CMB Anisotropy.

$$\frac{\Delta T(\vec{x}_o, \vec{n})}{T_o} = \underbrace{\frac{1}{3}(\Phi(\vec{x}_e) - \Phi(\vec{x}_o))}_{\text{Sachs-Wolfe}} + \underbrace{(\vec{v}(\vec{x}_e) - \vec{v}(\vec{x}_o))\vec{n}}_{\text{Doppler}} - \underbrace{\frac{2}{c^2} \int_e^o dx \frac{\partial \Phi(x\vec{n})}{\partial x}}_{\text{Integrated Sachs-Wolfe}} + \underbrace{\frac{\Delta T(\vec{x}_e)}{T_o}}_{\text{Intrinsic}}$$


### ♠ Intrinsic Fluctuations.

$$\rho_\gamma \sim a^{-4} \sim T^4 \quad \Longrightarrow \quad \delta_\gamma = \frac{\delta \rho_\gamma}{\rho_\gamma} = 4 \frac{\delta T}{T_o} \quad \Longrightarrow \quad \frac{\delta T}{T_o} = \frac{1}{4} \delta_\gamma$$

If the density of the radiation field varies across the sky, it will generate temperature anisotropies. The relation between the different energy densities defines the **Fluctuation Type**.



## Adiabatic Fluctuations.



The relative number density of particles is the same at every point for all particle species.

$$\delta = \frac{\delta n_B}{n_B} = \frac{\delta n_X}{n_X} = \frac{\delta n_\gamma}{n_\gamma}$$

Since  $a^{-1} \sim T$ ,  $n_B \sim a^{-3}$  and  $n_\gamma \sim a^{-4}$  then:


$$\delta_\gamma = \frac{4}{3}\delta_B; \quad \frac{\delta T}{T_0} = \frac{1}{4}\delta_\gamma = \frac{1}{3}\delta_B$$

This fluctuations are termed **Adiabatic** because the baryon/photon ratio does not change anywhere, i.e., there is no heat flux:

$$\delta \left( \frac{n_B}{n_\gamma} \right) = \frac{n_B}{n_\gamma} \left[ \frac{\delta n_B}{n_B} - \frac{\delta n_\gamma}{n_\gamma} \right] = 0$$



## Isocurvature Fluctuations.



The total energy density is constant. Fluctuations in the radiation field are compensated by all other matter components.

$$\left. \begin{array}{l} \rho_m = m_X n_X \\ \rho_\gamma = \sigma T^4 \end{array} \right\} \Rightarrow \delta\rho = m_X \delta n_X + 4\sigma T^3 \delta T = 0 \Rightarrow \rho_X \frac{\delta n_X}{n_X} + 4\rho_\gamma \frac{\delta T}{T_o} = 0 \Rightarrow$$

$$\frac{\delta T}{T_o} = -\frac{1}{4} \frac{\rho_X}{\rho_\gamma} \frac{\delta n_X}{n_X}$$

For modes outside the horizon, the behaviour is different in RD and in the MD regimes.

- In RD perturbations are isothermal:  $\rho_X \ll \rho_\gamma \Rightarrow (\delta T/T_o) \sim 0$
- In MD large anisotropies will develop:  $\rho_X \gg \rho_\gamma \Rightarrow (\delta T/T_o) \gg (\delta n_X/n_X)$

Inside the horizon, isocurvature perturbations evolve to become adiabatic.



## Evolution of a Radiative Field in an Expanding Universe.

Let  $f$  be the distribution function of a gas of particles. If there are no particle creation, in an expanding Universe relativistic and non-relativistic particles verify  $p \sim a^{-1}$ , and

$$f(\vec{x}, \vec{p}) = \frac{d^6 N}{d^3 x d^3 p} \sim \frac{const}{a^3 a^{-3}} \sim const.$$

**The distribution functions DO NOT GET DISTORTED with the expansion.**

♠ For a planckian field:

$$f(\vec{x}, \vec{p}) = f(E/T)$$

Since with the expansion the energy of the photon distribution changes, so must do the temperature.

$$E \sim a^{-1} \quad \Rightarrow \quad T \sim a^{-1}$$



## Dipole.

♣ The local motion of the observer produces a temperature anisotropy with a dipole pattern.

Let a radiative field be distributed uniformly on a volume for an observer  $O$ . For any observer  $O'$  that moves with velocity  $\vec{v}$  with respect to  $O$ , the energy  $E$  of any photon will experience a boost:


$$E' = E\gamma(1 - \vec{k}\vec{v}/c) = E\gamma(1 - \frac{v}{c} \cos \theta)$$

where  $\theta$  is the angle between the photon and the direction of motion of the observer  $O'$ .

♠ Lorentz transformations do not affect the distribution function (Landau & Lifshitz, v.II), so the change boost in energy corresponds to a change in temperature:

$$T'(\theta) = T_o\gamma(1 - \frac{v}{c} \cos \theta) \quad \Rightarrow \quad \frac{\Delta T}{T_o} = \frac{T'(\theta) - T_o}{T_o} = \gamma(1 - \frac{v}{c} \cos \theta) \approx \frac{v}{c} \cos \theta$$

Earth motion gives rise to a dipole pattern.



**Dipole Pattern**

The Fixsen et al (1994, ApJ 420, 445) measurement of the CMB dipole was:  $3.343 \pm 0.016$  mK (95% confidence level) with a direction  $(\alpha, \delta) = (168^\circ.9, -7^\circ.5)$  that in Galactic coordinates are  $(l, b) = (265^\circ.26, 48^\circ.74)$



## Anisotropies due to the Gravitational Potential.

♣ The variation of the gravitational potential at the LSS induces temperature anisotropies (Sachs-Wolfe effect).

$$\frac{\Delta T(\vec{x}_o, \vec{n}_e, t_o)}{T_o} = \frac{1}{3} \Phi(\vec{x}_e, t_o)$$

The potentials will be evaluated today.

♣ Let us compute one observable of the temperature field: the correlation function.

$$\langle \Delta T(\vec{x}) \cdot \Delta T(\vec{x}') \rangle = \frac{1}{9} \int \frac{d^3 k}{(2\pi)^3} \frac{d^3 k'}{(2\pi)^3} \langle \phi(k) \phi^*(k') \rangle e^{i(\vec{k}\vec{x} - \vec{k}'\vec{x}')}$$

♣ Poisson's equation gives a relation between matter power spectrum and gravitational potential:

$$\nabla^2 \phi(\vec{x}, t_o) = 4\pi G \bar{\rho}(t_o) \delta(\vec{x}, t_o) a^2(t_o) = \frac{3}{2} H_o^2 \Omega_m \delta(\vec{x}, t_o) \quad \Rightarrow \quad \phi(\vec{k}, t_o) = \frac{3}{2} H_o^2 \Omega_m \frac{\delta(\vec{k}, t_o)}{k^2}$$





After substituting the previous expression:

$$\langle \Delta T(\vec{x}) \cdot \Delta T(\vec{x}') \rangle = \frac{1}{4} H_o^4 \Omega_m^2 \int \frac{d^3 k}{(2\pi)^3} \frac{d^3 k'}{(2\pi)^3} \frac{\langle \delta(k) \delta^*(k') \rangle}{k^2 k'^2} e^{i(\vec{k}\vec{x} - \vec{k}'\vec{x}')}$$

Taking into account that  $\langle \delta^*(\vec{k}) \delta(\vec{k}') \rangle = (2\pi)^3 \delta^D(\vec{k} + \vec{k}') P(k)$  and the Rayleigh expansion of a plane wave:

$$e^{-i\vec{k}\hat{n}r} = 4\pi \sum_{lm} i^l j_l(kr) Y_{lm}^*(\Omega_{\hat{k}}) Y_{lm}(\Omega_{\hat{n}}),$$

we obtain

$$C(\theta) = \langle \Delta T(\vec{x}) \cdot \Delta T(\vec{x}') \rangle = \frac{1}{4\pi} \sum (2l+1) C_l P_l(\cos \theta); \quad C_l = \frac{H_o^4}{2\pi} \int_0^\infty k^2 dk \frac{P(k)}{k^4} j_l^2(kR_H)$$

where  $\theta$  is the angle between directions of observation  $x\hat{n}$  and  $x'\hat{n}'$  and  $R_H = 2cH_o^{-1}$ .



In the special case that the power spectrum behaves like a power law:  $P(k) = Ak^n$  we get:

$$C_l = \frac{AH_o^{n+3} \Gamma(3-n) \Gamma\left(\frac{2l+n-1}{2}\right)}{16 \Gamma^2\left(\frac{4-n}{2}\right) \Gamma\left(\frac{2l+5-n}{2}\right)}$$

In the special case of  $n = 1$  then

$$C_l = \frac{AH_o^4}{4\pi} \frac{1}{l(l+1)} = 6C_2 \frac{1}{l(l+1)}$$